

Cardioprotective Potential of *Quisqualis indica* Leaves; An *in silico* Docking Study

Mayank Kulshreshtha^{1,*}, Anuj Kumar Sharma¹, Devender Pathak², Pawan Pandey³, Varsha Snehi², Karuna Shanker Shukla³, Akash Ved⁴, Arpita Singh⁵, Neeraj Verma⁶, Anita Sing⁷, Manjul Pratap Singh⁸

¹Department of Pharmacology, Rajiv Academy for Pharmacy, Mathura, Uttar Pradesh, INDIA.

²Department of Pharmaceutical Chemistry, Rajiv Academy for Pharmacy, Mathura, Uttar Pradesh, INDIA.

³Department of Pharmaceutics, Rajiv Academy for Pharmacy, Mathura, Uttar Pradesh, INDIA.

⁴Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Dr. A.P.J. Abdul Kalam University, Lucknow, Uttar Pradesh, INDIA.

⁵Department of Pharmaceutics, Seth Vishambhar Nath Institute of Pharmacy, Barabanki, Lucknow, Uttar Pradesh, INDIA.

⁶Department of Pharmacology, Hygia College of Pharmacy, Lucknow, Uttar Pradesh, INDIA.

⁷Department of Pharmaceutics, Sundeeep Pharmacy College, Ghaziabad, Uttar Pradesh, INDIA.

⁸Department of Pharmaceutics, ABESIT College of Pharmacy, Ghaziabad, Uttar Pradesh, INDIA.

ABSTRACT

Background: Nowadays cardiac problems are the main cause of death. It includes coronary artery disease, angina pectoris (stable and unstable), congestive heart failure, etc. **Objectives:** The aim of this paper was to generate scientific data regarding *in silico* analysis of beta-sitosterol, coumaric acid, lupeol, quercetin, and urosolic acid of Muscarinic (M₂) receptor. **Materials and Methods:** The leaf powder was treated with different reagents and prepared with different extracts. The phytochemical screening was carried out by treated the extracts with different reagents for the presence of various metabolites. RCSB protein data bank had used for docking studies. **Results:** The phytochemical screening clearly revealed the presence of various metabolites like flavonoids, alkaloids, saponins, etc. *In-silico* analysis of beta-sitosterol, coumaric acid, lupeol, quercetin, and urosolic acid had very good interactions with cholinergic receptor (M₂). The obtained score is -7.96, -5.63, -6.7, -7.73, -5.82 for beta-sitosterol, coumaric acid, lupeol, quercetin, and urosolic acid, respectively, which lies in the standard scale. **Conclusion:** All these metabolites (compounds) are present in *Quisqualis indica* leaf extracts (aqueous and ethanolic) and found to be good cardio protective agents.

Keywords: *Quisqualis indica*, Cardio-protective, *in silico*.

Correspondence:

Dr. Mayank Kulshreshtha

Professor, Rajiv Academy for Pharmacy,
Mathura, Uttar Pradesh, INDIA.

Email: professorsmayank@gmail.com

Received: 14-03-2023;

Revised: 24-03-2023;

Accepted: 02-04-2023.

INTRODUCTION

Depending on a patient's physical makeup or a disease, the cardiac system may work differently from one patient to the next, and breathing may cause subtle variances. In extremely rare instances, the complete human structure may resemble a mirror image while still being normal or when a congenital heart abnormality is present. The body structures that typically exhibit lateralization are organised in isomeric way in various situations, which are more frequent than the mirror-imaged condition but still very uncommon.¹ The heart of a person is a crucial organ that continuously pumps blood in cycles throughout the body. The diaphragm, which is close to the thoracic cavity, supports the heart. It is located in the mediastinum, an area of the body where the sternum and spinal column meet anatomically (rib to the

diaphragm, and between the lungs). The heart is always around two-thirds to the left of the body's midline. The pericardium encircles and shields the heart. It aids in keeping the heart confined to its location in the mediastinum and provides enough room for forceful, quick contraction. The epicardium (external layer), the myocardium (middle layer), and the endocardium are the three layers that make up the heart's wall (inner layer). Two inferior pumping chambers and two superior receiving chambers make up the heart (known as ventricles).²

Quisqualis indica Linn. (*Q. indica*) belongs to family-*Combretaceae* can reach up to 9 meters. Locally known as 'Malti' It is cultivated all parts of India but also find in Africa and Indo-Malaysian region.^{3,4} It generally requires an area with full sunlight and regular watering.³ Leaf and root extracts/juices are traditionally used as anthelmintics, and leaves are used to ease flatulence. Externally, a leaf infusion is used to cure boils and ulcers. To expel worms, seeds are anthelmintic to youngsters.^{4,5} It exhibits pharmacological activities such as excitotoxicity,^{6,7} anti-inflammatory,^{8,9} antipyretic,¹⁰ immunomodulatory,^{11,12}



DOI: 10.5530/fra.2023.1.7

Copyright Information :

Copyright Author (s) 2023 Distributed under
Creative Commons CC-BY 4.0

Publishing Partner : EManuscript Tech. [www.emanuscript.in]

anti-staphylococcal activity,¹³ acetyl cholinesterase inhibitor,^{14,15} antioxidant activity,¹⁶ and so on. This plant has a number of medicinally active phytochemical components that are responsible for a variety of pharmacological effects.

According to the literature review, there is no research-based data on *in silico* docking investigations on the leaves of *Q. indica*. As a result, the current research was carried out to evaluate the above-mentioned potential of leaves using WHO criteria and other official procedures. For the first time, a method for determining the quality of several metabolites in *Q. indica* has been devised that is relevant, responsive, and predictable. This research data may be used as a reference for new research, plant physiology, chemistry analysis, biochemistry, and the preparation of a plant monograph.

MATERIALS AND METHODS

Collection and authentication of leaf material

The leaves of *Q. indica* were collected from National Botanical Research Institute, Lucknow, India and authenticated by National Institute of Science Communication and Information Resources, Delhi, India (NISCAIR/RHMD/Consult/2015/2862/55-1).

Drugs and chemicals

Ethanol was purchased from Changshu Yangyuan Chemical, China, and the reagents had used in phytochemical screening, they were freshly prepared. All chemicals were analytical grade.

Extraction procedure

Leaves were dried on cool place and powdered with the help of electric grinder. Soxhlet method was followed in which powder was defatted using 250 mL of petroleum ether for more than 6 hr, after that the powders were dried and extracted with different solvents. The obtained final fractions were concentrated under vacuum in a rotary evaporator at 40°C and stored at 4°C for further use.¹⁷

Phytochemical screening

Different metabolites like carbohydrates, alkaloids, flavonoids, tannins, proteins, terpenoids etc present in different leaf extracts of *Q. indica*.

In silico docking studies

Molecular docking of beta-sitosterol, coumaric acid, lupeol, quercetin and urosolic acid to muscarinic (M₂) receptors: To predict the contact of the drug molecule with receptor, the molecular docking simulation was carried out.¹⁸ It involved the following steps.

Selection and preparation of protein

The RCSB protein data library was used to download a muscarinic (M₂) receptor coupled to ligand (PDB: 5ZHP).

Preparation of ligands

The ligand Tiotropium was prepared for molecular docking simulation by using the Auto Dock software to determine the number of rotatable, non-rotatable, and unrotatable bonds present in the ligand.

Grid box formation

The grid parameter points of the grid box required to perform the molecular docking simulation of ligand molecules were enumerated using the receptor binding site, which was found using the protein visualisation software PyMol. For all docking runs, these grid parameters were used. To ensure that all extended conformations of the ligand fit within the grid box, it was put in the middle of the ligand by covering all binding residues involved in the binding of the ligand.

Grid map preparation

The map files for different atom types in ligands and receptor viz. A, C, F, HD, N, OA, and SA were prepared by running Auto grid utility of the Auto dock suite.

Docking parameters

Lamarckian Genetic Algorithm (LGA) is the primary conformational search approach used in Auto Dock for molecular docking simulation. Docking parameter file for each ligand was prepared by using 150 Genetic Algorithm (GA) runs, 250000 maximum numbers of evaluations, 27000 maximum number of generations, and 0.02% rate of gene mutation.

Docking method validation






The position and orientation of the ligand obtained after the molecular docking study represent the potential binding modes of the inhibitors. The various docking parameters considered in the docking methods were validated by redocking individually crystallized ligands beta-sitosterol, coumaric acid, lupeol, quercetin and urosolic acid over the muscarinic acetylcholine

Table 1: Phytochemical analysis of different extracts of *Q. indica*.

Phytoconstituents	AEQI	EEQI
Alkaloids	Present	Present
Glycosides	Present	Present
Tannins	Present	Present
Flavonoids	Present	Present
Fats and oil	Present	Present
Carbohydrates	Present	Present
Reducing sugar	Present	Present
Proteins	Present	Present
Saponin	Absent	Absent
Terpenoids	Present	Present

AEQI- Aqueous extract of *Q. indica*, EEQI- Ethanolic extract of *Q. indica*

Table 2: Docking results of certain drugs with M₂ Receptor (3UON).

Sl. No.	Name of drugs	Structure	Binding energy (Kcal/mol)
1	Beta-sitosterol		-7.96
2	Coumaric acid		-5.63
3	Lupeol		-6.7
4	Quercetin		-7.73
5	Urosolic acid		-5.92

receptor (M₂). The molecular docking simulation technique is validated by using different parameters.

RESULTS

Phytochemical testing of different extracts shows the presence of alkaloids, glycosides, sterols, carbohydrates, tannins, terpenoids, saponins and flavonoids as shown in Table 1 respectively. From Table 2, the maximum binding energy was obtained -7.96 Kcal/mol in case of beta-sitosterol.

An appropriate grid box was prepared by covering all residues involved in the active binding of the ligand 9EC to the M₂ muscarinic acetylcholine receptor. The coordinates used for the preparation of the grid box are tabulated in Table 3. Three-dimensional grid box covering and the active ligand binding sites present in the receptor molecule as well as all residues involved in the binding of the ligand show in Figure 1. Molecular Docking results of ligands with M₂ muscarinic acetylcholine receptor are shown in Table 4.

In case of validation, the following parameters were used to validate the molecular docking process for docking of muscarinic acetylcholine receptors with different legends.

Binding energy

Molecular docking should fall in the normal range of -5 to -15 kcal/mol. The molecular docking validated as the binding energy of the ligands beta-sitosterol, coumaric acid, lupeol, quercetin and urosolic acid with the M₂ muscarinic acetylcholine receptor was found to be -7.96, -5.63, -6.7, -7.73 and -5.82 kcal/mol respectively, which lies in the predefined range of -5 to -15 kcal/mol.

Overlay methods

The overlaid conformation of the docked ligand with reference to the bioactive crystal structure of the ligand was obtained from RSCB protein data bank.

Chemical resemblance


The molecular docking method is validated when the docked ligand should have the same interactions with the residues of the macromolecule as those present in the downloaded crystallized macromolecule. The interactions present in the crystal structure and the docked structure are shown in Figure 2.

Table 3: Coordinates of the grid box for the muscarinic acetylcholine receptor (M₂).

Proteins	x-D	y-D	z-D	Spacing (Å)	x center	y center	z center
5ZHP	40	40	40	0.375	8.15	0.745	-3.75


Table 4: Molecular Docking results of ligands with M₂ muscarinic acetylcholine receptor.

Proteins	Interacting residues	Binding energy (kcal/mol)
5ZHP	Beta-sitosterol	-7.96
	Coumaric acid	-5.63
	Lupeol	-6.7
	Quercetin	-7.73
	Ursolic acid	-5.82

**Figure 1:** Three dimensional grid box covering the active ligand binding sites present in the receptor molecule as well as all residues involved in the binding of the ligand.

DISCUSSION

Cardiovascular diseases are the main and most important cause of death. According to Nature Reviews, Cardiology, Cardiovascular diseases are also the leading cause of death in mostly countries like China etc.¹⁸ There are lots of types of diseases and they are all dangerous for heart in which coronary artery diseases, and also atherosclerotic heart disease, is the condition in which the accumulation of plaque within the walls of coronary arteries. Coronary artery disease is the leading cause of death worldwide. While the symptoms and signs of coronary artery disease are

**Figure 2:** Binding mode and chemical interactions of the bound ligands (1) Beta-sitosterol, (2) Coumaric acid, (3) Lupeol, (4) Quercetin, and (5) Ursolic acid within the active ligand binding site of M₂ muscarinic acetylcholine receptor.

noted in the advanced state of disease, most individuals with coronary artery disease show no evidence of disease for decades as the disease progresses before the first onset of symptoms, often a "sudden" heart attack, finally arises.^{19,20} Cholinergic receptors are classified into muscarinic and nicotinic, in which muscarinic is again classified into M₁, M₂, M₃, M₄, and M₅. The M₂ receptor has been selected for docking studies because the main location of M₂ is heart. Beta-sitosterol, coumaric acid, lupeol, quercetin and ursolic acid have been selected for docking studies because they all have a protective mechanism of heart.²¹ The 5ZHP protein complex of RCSB protein data bank consists of four identical chains of amino acids and bound ligand 9EC [(1R,2R,4S,5S,7s)-7-({[4-fluoro-2-(thiophen-2-yl)phenyl]carbamoyl}oxy)-9,9-dimethyl-3-oxa-9-azatricyclo[3.3.1.0~2,4~]nonan-9-ium]. After processing the receptor molecule it was saved in "pdbqt" format by using AutoDock software. Beta-sitosterol is widespread used as a lipid lowering agent and improves the condition of prostate gland. Beta-sitosterol may reduce the risk of Cardiovascular Disease (CVD) by lowering LDL cholesterol in the blood.²² It is clearly proved that beta-sitosterol is a good cardioprotective agent. Phytosterols like beta-sitosterol are homologues to cholesterol found in various fruits and vegetables and present in the daily

diet as similar to cholesterol (200–400 mg/kg).²³ Beta-sitosterol has a lipid lowering quality because of competitive inhibition of cholesterol absorption and a gene implicated in the metabolism of cholesterol.²⁴ Coumaric acid is a well-known phenolic compound that is also present in various fruits and vegetables and reported to have anti-inflammatory activity. p-coumaric acid can convert to phenolic acids such as chlorogenic acid, rosmarinic acid, flavonoids, and other secondary metabolites and possesses various effects including antioxidant, anti-angiogenic, anti-UV damage, and antiplatelet properties.^{25,26}

Lupeol has also reported various pharmacological properties like antioxidant, anti-inflammatory, anti-hyperglycemic, anti-dyslipidemic, and anti-mutagenic effects.²⁷ Quercetin exhibits significant heart-related benefits such as inhibition of LDL oxidation, endothelium-independent vasodilator effects etc. Urosolic acid also has the good score and it exerts these effects in various tissues and organs. Mechanisms include suppressing nuclear factor-kappa B signaling in cancer cells, improving insulin signaling in adipose tissue, reducing the expression of markers of cardiac damage in the heart.²⁸

CONCLUSION

Looking to future public health approaches, all these compounds are targeted for cardiovascular disease prevention. This paper has clearly emphasized the *Q. indica* has good cardioprotective potential, but detailed research is needed for prospects.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ABBREVIATIONS

Q. indica: *Quisqualis indica*; **M₂:** Muscarinic receptor; **AEQI:** Aqueous extract of *Q. indica*; **EEQI:** Ethanolic extract of *Q. indica*.

REFERENCES

- Anderson RH, Webb S, Brown NA. Defective lateralisation in children with congenitally malformed hearts. *Cardiol Young*. 1998;8(4):512-31. doi: 10.1017/s1047951100007204, PMID 9855108.
- Tortora G, Derrickson B. Principles of anatomy and physiology. 14th ed. Wiley; 2009. p. 718-21.
- Joshi SG. Medicinal plants. 1st ed. Oxford and IBH Publishing Company; 2002. p. 11-2.
- Shih CT. 'Stuartexchange' niyog niyogan, Art guild for education and communication foundation Inc; 2011.
- Kirtikar KR, Basu BD. Indian Medicinal plant. 2nd ed. Asistic Publishing House; 2006. p. 14-6.

- Munir M, Lu L, Mcgonigle P. Excitotoxic cell death and delayed rescue in human neurons derived from NT2 cells. *J Neurosci*. 1995;15(12):7847-60. doi: 10.1523/JNEUROSCI.15-12-07847.1995, PMID 8613724.
- Murphy TH, Baraban JM. Glutamate toxicity in immature cortical neurons precedes development of glutamate receptor currents. *Brain Res Dev Brain Res*. 1990;57(1):146-50. doi: 10.1016/0165-3806(90)90195-5, PMID 1982523.
- Yashraj Y, Mohanty PK, Kasture SB. Anti-inflammatory activity of hydroalcoholic extract of *Quisqualis indica* Linn. Flower in rats. *Int J Pharm Life Sci*. 2011;2:977-81.
- Gautam R, Jachak MS. Naturally occurring polyphenols with anti-inflammatory activity. *CRIPS*. 2007;8(4):20-32.
- Nitu S, Pankaj K, Samantha KC, Reena D. Antipyretic activity of methanolic extract of leaves of *Quisqualis indica* Linn. *IJPRD*. 2010;2(9):122-6.
- Sagrawat H, Khan Y. Immunomodulatory plants: A phytopharmacological review. *Pharmacogn Rev*. 2007;1(2):248-58.
- Yadav Y, Mohanty PK, Kasture SB. Evaluation of immunomodulatory activity of hydroalcoholic extract of *Quisqualis indica* Linn. flowers in Wistar rats. *IJPLS*. 2011;2:946-7126.
- Jahan FN, Rahman MS, Mukhlesur R, Simon G, Mohammad MM, Samir KS. Diphenylpropanoids from *Quisqualis indica* Linn. and their anti-staphylococcal activity. *Lat Am J Pharm*. 2009;28(2):279-83.
- Wetwitayaklung P, Limmatvapirat C, Phaechamud T, Keokitichai S. Kinetics of acetylcholinesterase Inhibition of *Quisqualis indica* Linn. flower extract. *Silpakorn Sci Technol J*. 2007;1:20-8.
- Kaisar MA, Islam MR, Rahman MS, Hossain MK, Rashid MA. Total phenolic content, free radical scavenging activity and reducing power of *Quisqualis indica* Linn. *Dhaka Univ J Pharm Sci*. 2009;8(2):173-5. doi: 10.3329/dujps.v8i2.6034.
- Solanki R, Gupta A, Tripathy A, Soni D, Jana G. Pharmacognostic evaluation of the leaf of *Rhus succedanea*. *J Nat Prod Plant Resour*. 2011;1(4):20-6.
- Jie G, Peng Z, Xiao-qin Z, Xue-feng Z, Guang-yao W. Computational methods for designing potential inhibitors for activin type IIb (ActRIIB) receptor for treatment of anaemia. *Bio Resh*. 2017;28(8):3369-76.
- Zhao D, Liu J, Wang M, Zhang X, Zhou M. Epidemiology of cardiovascular disease in China: current features and implications. *Nat Rev Cardiol*. 2019;16(4):203-12. doi: 10.1038/s41569-018-0119-4, PMID 30467329.
- Benjamin EJ, Muntner P, Alonso A, Bittencourt MS, Callaway CW, Carson AP, et al. Heart disease and stroke statistics—2019 update: A report from the American Heart Association. *Circulation*. 2019;139(10):e56-e528. doi: 10.1161/CIR.0000000000000659, PMID 30700139.
- [cited Jan 04 2021] Available from: <https://medicalxpress.com/conditions/coronary-artery-disease>.
- Hacker M, Bachmann K, Messer W. Pharmacology – principles and practices. 1st ed. Elsevier Publication; 2009. p. 112-6.
- Racette SB, Lin X, Lefevre M, Spearie CA, Most MM, Ma L, et al. Dose effects of dietary phytosterols on cholesterol metabolism: a controlled feeding study. *Am J Clin Nutr*. 2010;91(1):32-8. doi: 10.3945/ajcn.2009.28070, PMID 19889819.
- Ostlund RE. Phytosterols in human nutrition. *Annu Rev Nutr*. 2002;22(1):533-49. doi: 10.1146/annurev.nutr.22.020702.075220, PMID 12055357.
- Calpe-Berdiel L, Escolà-Gil JC, Blanco-Vaca F. New insights into the molecular actions of plant sterols and stanols in cholesterol metabolism. *Atherosclerosis*. 2009;203(1):18-31. doi: 10.1016/j.atherosclerosis.2008.06.026.
- Yoon JH, Youn K, Ho CT, Karwe MV, Jeong WS, Jun M. p-Coumaric acid and ursolic acid from corni fructus attenuated β -Amyloid25–35-induced toxicity through regulation of the NF- κ B signaling pathway in PC12 cells. *J Agric Food Chem*. 2014;62(21):4911-6. doi: 10.1021/jf501314g, PMID 24815946.
- Khairy M, Dianat M, Badavi M, Mard SA, Bayati V. p-Coumaric acid protects cardiac function against lipopolysaccharide-induced acute lung injury by attenuation of oxidative stress. *Iran J Basic Med Sci*. 2019;22(8):949-55. doi: 10.22038/ijbms.2019.36316.8650, PMID 31579452.
- Tsai FS, Lin LW, Wu CR. Lupeol and its role in chronic diseases. *Adv Exp Med Biol*. 2016;929:145-75. doi: 10.1007/978-3-319-41342-6_7, PMID 27771924.
- Seo DY, Lee SR, Heo JW, Rhee BD, Ko KS, et al. Ursolic acid in health and disease. *Korean J Physiol Pharmacol*. 2018;22(3):235-48. doi: 10.4196/kjpp.2018.22.3.235, PMID 29719446.

Cite this article: Kulshreshtha M, Sharma AK, Pathak D, Pandey P, Snehi V, Shukla KS, et al. Cardioprotective potential of *Quisqualis indica* leaves; An *in silico* docking study. *Free Radicals and Antioxidants*. 2023;13(1):41-5.